

A survey of UV-excess AGNs in the South Galactic Pole^{*}

A sample for the analysis of the QSO clustering

F. La Franca¹, C. Lissandrini², S. Cristiani^{2,3}, L. Miller⁴, M.R.S. Hawkins⁵, and H.T. McGillivray⁵

¹ Dipartimento di Fisica, Università degli studi Roma Tre, Via della Vasca Navale 84, I-00146 Roma, Italy

² Dipartimento di Astronomia, Università di Padova, Vicolo dell'Osservatorio 5, I-35122, Padova, Italy

³ Space Telescope-European Coordinating Facility, European Southern Observatory, Karl-Schwarzschild Str. 2, D-85748 Garching b. München, Germany

⁴ Department of Physics, Nuclear and Astrophysics Laboratory, Keble Road, Oxford OX1 3RH, UK

⁵ Royal Observatory of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK

Received July 8, 1999; accepted ???? ??, 1999

Abstract. Spectra, position, magnitudes and colors are presented for 485 faint ($B_J < 20.5$) emission line objects selected with the ultraviolet-excess (UVX) criterion on a area of 24.6 deg^2 in the South Galactic Pole. The objects were selected from the analysis of pixel-to-pixel stacking of COSMOS scans of UKST U J and R plates. The candidates were observed with the Meudon-ESO Fiber Optics System (MEFOS) at the ESO 3.6m telescope. 429 type 1 AGNs have been identified (373 in the redshift range $0.3 < z \leq 2.2$). This sample has allowed the measure of a difference on the QSO clustering evolution in comparison with that found for galaxies (La Franca et al. 1998). The region is part of the ESO Imaging Survey (EIS) and of the 2dF QSO redshift survey.

Key words: Catalogs – Surveys – quasars: general

1. Introduction

In the last decade the study of the evolution of the QSO luminosity function and of the clustering properties of QSOs has been based on statistical analysis of even larger QSO catalogues. For redshifts lower than 2.2 the color techniques provide good selection methods. Now days the QSO luminosity function evolution at $z < 2.2$ is well established (e.g. La Franca & Cristiani 1997, Goldschmidt & Miller 1998). This kind of studies are mainly based on the most statistically significant QSO samples such as the Durham/AAT QSO sample (Boyle et al. 1990) at faint magnitudes ($B < 20.9$), and the Bright QSO Survey (Schmidt & Green 1983), the Large Bright QSO Survey

(Hewett et al. 1995), the Edinburgh UVX QSO survey (Goldschmidt et al. 1992, Miller et al. in prep), the Homogeneous Bright QSO Survey (HBQS, Cristiani et al. 1995) at bright magnitudes ($B < 18.5$). In total these samples collect about a thousand of QSOs with $z < 2.2$. But these samples are still not ideally suited to study the QSO clustering and its evolution (see Andreani & Cristiani 1992). The best QSO samples for clustering analysis should have the highest possible surface densities and cover a contiguous areas, such as the planned QSO 2dF redshift survey (Croom et al. 1998).

In order to improve the signal-to-noise ratio in the estimate of the clustering of QSOs, we have built a new sample of 429 QSOs down to $B_J = 20.5$ over a contiguous area of 24.6 deg^2 . This sample has allowed the measure of a difference on the QSO clustering evolution in comparison with that found for galaxies (La Franca et al. 1998).

The survey is located in the south Galactic pole (SGP) region, where some of the Durham/AAT sample areas are included (Boyle et al. 1990), and part of the high-redshift QSO survey of Warren et al. (1991b) was carried out. The region has been also studied by Campusano (1991). The central area of 1.7 deg^2 in our region has been covered by the ESO Imaging Survey in the B, V and I band (EIS, da Costa 1998, Nonino et al. 1999, Prandoni et al. 1999, Zaggia et al. 1999). The acquisition of the corresponding U data is planned for 1999. Further deep multicolor imaging over an area of about half a sq.deg. is planned with the WFI at the ESO 2.2m telescope. Moreover about 10 deg^2 of the southern area of the region is included in the 2dF QSO redshift survey (Croom et al. 1998).

2. The construction of the catalogue

Our survey is based on a set of UKST plates in the U, B_J and R bandpass (see Blair & Gilmore (1982) for a definition of the various systems). UKST plates subtend $6^\circ.4 \times 6^\circ.4$. Details of the photographic plate material are

Send offprint requests to: F. La Franca

* Based on material collected with the UK Schmidt Telescope, with the ESO-La Silla Telescopes and on COSMOS scans.

Correspondence to: lafranca@fis.uniroma3.it

Table 1. UKST photographic plates

emulsion	filter	number	date	exp. time (min.)
IIaO	UG1	U 2639	1976-09-28	180
IIIaJ	UG1	U 6326	1980-09-03	180
IIIaJ	UG1	U 6380	1980-09-15	180
IIIaJ	GG395	J 9764	1984-11-23	70
IIIaJ	GG395	J 9765	1984-11-23	70
IIIaJ	GG395	J 9766	1984-11-23	70
IIIaJ	GG395	J 9770	1984-11-24	70
IIIaJ	GG395	J 9771	1984-11-24	70
127-04	RG630	R 3498	1977-08-15	90
IIIa-F	RG630	R 4676	1978-12-02	90
IIIa-F	RG630	R 9594	1984-09-22	90
IIIa-F	RG630	R 9672	1984-10-17	90
IIIa-F	RG630	R11331	1986-09-02	90

listed in Table 1. The plates have been scanned in Mapping and Imaging Mode on the COSMOS microdensitometer (MacGillivray & Stobie 1984). The mapping mode used a spot size of $16 \mu\text{m}$ (FWHM), corresponding to 1 arc-sec. For each band-pass the digitized Mapping Mode data have been added together in order to increase the signal to noise ratio. The reliability of the co-adding technique has been verified by many tests on UKST plates (Hawkins 1991). The resulting coadded digitized data has been analyzed by the COSMOS crowded-field analysis software (Beard et al. 1990). The resulting tables contained the instrumental magnitudes, the area above the threshold, the intensity weighted x and y positions, and other useful parameters. We discriminated between point-like and extended sources using the COSMOS image parameters, that is, on morphological grounds. We defined a plane in which stellar locus was well determined, and drew the separation line near the locus. For typically $B_J < 19.5$, we used the $\log(\text{isophotal area})$ versus magnitude plane. At fainter magnitudes, where stellar peak surface brightness (I_{peak}) are not saturated, the $\log(I_{\text{peak}} - I_{\text{sky}})/I_{\text{sky}}$ versus magnitude plane provided a better separation. The two separation lines produced the same star-galaxy ratio at the overlap magnitude. Relative photometry, using instrumental magnitudes, was produced by COSMOS. Calibration was based on published sequences (Hawkins & Bessell 1988, Warren et al. 1991a).

The candidates were extracted from a rectangle in the sky centered at $\alpha(1950): 00^{\text{h}} 50^{\text{m}} 34.3^{\text{s}}$ and $\delta(1950): -28^{\circ} 10' 08''$ with limits $\Delta\alpha_{\text{min}} = -2.170^{\circ}$, $\Delta\alpha_{\text{max}} = 2.715^{\circ}$, $\Delta\delta_{\text{min}} = -2.520^{\circ}$, $\Delta\delta_{\text{max}} = 2.520^{\circ}$ (i.e. the limits in right ascension are the two lines distant $\Delta\alpha$ from the meridian at $\alpha(1950): 00^{\text{h}} 50^{\text{m}} 34.3^{\text{s}}$; see Cristiani et al. 1995, their Table 1). All the objects inside a radius of 0.15 degrees centered at $\alpha(1950): 00^{\text{h}} 50^{\text{m}} 17.43^{\text{s}}$, and $\delta(1950): -26^{\circ}$

Table 2. The observing log

Date	Number of fields	Integration time (min)
10-Oct-93/15-Oct-93	15	40 – 70 × 2
30-Oct-94/02-Nov-94	14	45 – 60
22-Sep-95/25-Sep-95	8	40 – 50
22-Nov-95/25-Nov-95	8	60 – 80

$51' 25.2''$ have been excluded from the catalogue as this region includes a crowded globular cluster. It results a total area of 24.55 deg^2 .

We have selected as candidates all the UVx "not extremely extended" objects with $B_J \leq 20.5$, satisfying a type of modified Braccisi less-restricted two color criterion (La Franca et al. 1992, Cristiani et al. 1995). The completeness of the selection has been tested against the QSOs already known in the field with redshift in the range $0.3 < z \leq 2.2$. We have selected 92% of the 176 blue QSOs already known in the field with $B_J \leq 20.5$. The completeness becomes 89% if the "red" QSOS from Warren et al. (1991a) are included in the comparison.

3. The spectroscopic survey

We have used the 3.6m ESO telescope, equipped with the Meudon-ESO Fiber Optic System MEFOS (Bellenger et al. 1991), to obtain low-resolution spectra of the QSO candidates in four observational campaigns between 1993 and 1995. MEFOS is a multifibre positioner which enables the conventional use of the ESO Boller & Chivens Spectrograph to be extended to multiobject spectroscopy. The spectra of up to 29 objects inside the 1° field of the prime focus of the 3.6m telescope can be simultaneously recorded. It consists of 29 remotely controlled arms, each carrying two *spectroscopic fibers* of 2.5 arcsec of diameter, one for the object, and one for the sky sampling. We used the ESO grating n. 13 (150 grooves/mm) with a resolution of $35 \text{ \AA}$ in the wavelength range $3600 - 8400 \text{ \AA}$. The CCD was a 512×512 Tektronics with $27 \mu\text{m}$ pixels (ESO n. 32). Wavelength calibration was carried out by comparison with exposures of He and Ar lamps. The campaign dates with the number of pointings and the exposure time are listed in Table 2. The data reduction was performed under MIDAS and followed the optimized spectra extraction from fiber spectrographs of Lissandrini et al. (1994). No absolute flux calibration has been applied. The spectra have been flux calibrated in relative fluxes just for the sake of facilitating the identification. Only the most evident cosmic rays hits have been removed.

4. The catalogue

Altogether 769 QSOs candidates have been identified (502 from the MEFOS campaign), 485 of which actually turned out to be emission line extragalactic objects. In Fig. 1 the

spectra of all the newly identified objects are shown. The S/N of the spectra permitted to recognize emission-line objects without leaving much doubt of misidentification. Only 15 objects have a dubious redshift identification. In Table 3 the complete list of the 485 emission line objects and their spectroscopic identification is given. The identification classes are: QSO for broad emission line type 1 AGNs (QSOs and Seyfert 1 galaxies); NL for narrow emission line galaxies (such as Seyfert 2, LINERS, Starburst or HII emissions). The uncertainty in the redshift estimation is of 0.001 rms. At magnitudes brighter than $B < 18.7$ the sample is complete and the QSOs are included in the catalogue of the Homogeneous Bright QSO Survey (HBQS, Cristiani et al. 1995). A “d” in the comments identify the objects for which the redshift identification is dubious. A total of 429 type 1 AGNs have been identified, 373 in the redshift range $0.3 < z \leq 2.2$. The photometry has an uncertainty of about 0.1 mag, while the astrometry has an accuracy of 1'' rms.

MEFOS has been decommissioned by ESO in 1995, and consequently at magnitudes fainter than $B=18.7$ a fraction of the QSO candidates has not been spectroscopically identified. The fainter sample is not useful for statistical applications aimed at measuring the evolution of the luminosity function of AGNs. However, the sample can be used for analysis of the clustering of QSOs by using the technique of scrambling the redshift distribution in the generation of the “random” data set used in the computation of the correlation function (see La Franca et al. 1998).

Table 3. List of the Candidates

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 40 33.0	-30 24 10	17.93	-0.89	+0.09	QSO	0.609	a
00 40 41.1	-29 17 22	17.75	-0.89	+0.17	QSO	2.084	b
00 40 46.0	-29 19 39	18.37	-0.77	+0.06	QSO	0.624	a
00 40 50.7	-30 18 01	18.34	-1.04	+0.19	QSO	0.496	b
00 40 54.8	-26 55 30	17.34	-0.52	+0.06	QSO	1.002	b
00 41 10.4	-26 12 47	19.72	-0.41	-0.08	QSO	1.720	c
00 41 14.9	-26 38 33	18.24	-0.54	+0.37	QSO	3.053	a
00 41 23.9	-28 44 06	18.18	-0.27	+0.15	QSO	0.839	a
00 41 28.4	-29 04 16	17.80	-0.61	-0.03	QSO	0.674	a
00 41 30.8	-26 07 38	17.30	-0.78	+0.36	QSO	2.501	a
00 41 38.1	-26 58 27	18.54	-0.47	+0.34	QSO	2.457	a
00 41 39.1	-27 54 32	20.03	-0.46	+0.32	QSO	1.112	
00 41 40.6	-28 59 36	17.97	-0.80	+0.12	QSO	2.134	a
00 41 42.4	-27 48 03	19.80	-1.26	+0.61	QSO	1.384	
00 41 43.1	-27 28 20	18.81	-1.00	+0.96	QSO	0.381	
00 42 07.2	-29 08 15	18.29	-1.03	+0.35	QSO	1.242	a
00 42 16.9	-25 50 32	18.27	-0.38	+0.15	QSO	0.454	a
00 42 18.8	-27 30 28	19.66	-0.82	+0.80	NL	0.287	
00 42 20.6	-27 12 05	19.17	-1.48	+0.54	QSO	1.275	
00 42 20.7	-28 25 46	18.80	-0.53	+0.14	QSO	0.567	
00 42 26.6	-27 50 19	18.13	-0.74	-0.10	QSO	0.741	a
00 42 40.9	-27 44 40	20.24	-0.73	+0.84	NL	0.444	
00 42 41.7	-29 30 58	17.92	-0.48	+0.40	QSO	2.388	a
00 42 51.0	-27 31 48	20.28	-1.15	+0.24	QSO	1.927	
00 42 54.7	-27 39 38	19.27	-0.35	+0.15	QSO	2.430	c
00 42 55.0	-27 29 05	18.73	-0.70	+0.53	QSO	0.933	a
00 42 55.6	-29 15 23	19.18	-0.53	+0.41	NL	0.206	
00 43 01.8	-29 37 03	18.77	-0.57	+0.05	QSO	2.240	c
00 43 02.6	-27 44 13	19.81	-0.78	+0.18	QSO	1.760	
00 43 15.0	-30 30 49	19.13	-0.54	+0.17	QSO	2.390	c
00 43 22.3	-27 43 32	18.46	-0.70	+0.35	QSO	1.049	a
00 43 22.7	-27 42 15	19.96	-0.70	+1.29	NL	0.381	d
00 43 31.1	-28 19 52	20.02	-0.20	+0.25	NL	0.050	
00 43 32.4	-28 45 36	19.64	-1.18	+0.36	QSO	2.180	e
00 43 38.2	-26 44 28	19.28	-1.52	+0.87	QSO	1.041	a
00 43 40.3	-27 03 01	20.11	-0.58	+0.60	NL	0.111	
00 43 49.7	-30 06 36	18.65	-1.00	+0.37	QSO	1.124	a
00 43 53.6	-27 55 12	20.08	-1.10	+0.05	QSO	1.907	
00 43 55.2	-30 29 26	18.72	-0.59	+0.25	QSO	2.190	e
00 43 57.9	-28 01 41	19.84	-1.28	+0.77	QSO	1.591	
00 44 00.7	-27 01 06	19.57	-0.78	+0.64	QSO	2.166	
00 44 04.8	-28 43 01	19.62	-0.26	+0.25	QSO	2.230	
00 44 10.5	-27 13 49	20.27	-1.18	+0.10	QSO	2.168	
00 44 10.6	-26 11 25	17.52	-0.77	+0.52	QSO	0.129	a
00 44 18.1	-27 38 54	20.13	-0.49	+1.03	NL	0.330	
00 44 21.1	-26 58 09	19.20	-0.97	+0.38	QSO	0.968	
00 44 22.5	-29 51 22	18.12	-0.44	+0.48	QSO	0.207	a
00 44 30.0	-28 28 26	20.41	-0.43	+0.24	NL	0.186	
00 44 48.6	-28 49 01	19.36	-0.37	+0.01	QSO	1.887	
00 44 52.4	-28 16 22	20.05	-0.75	+0.35	QSO	1.786	
00 44 52.7	-26 19 21	20.17	-0.38	+0.27	NL	0.073	
00 44 53.9	-27 32 30	20.38	-1.41	+0.57	QSO	0.639	
00 44 55.0	-26 09 40	20.45	-1.01	-0.21	QSO	1.911	
00 44 56.8	-26 52 30	20.33	-0.52	+0.35	QSO	1.078	
00 44 57.5	-26 33 31	19.48	-0.76	+0.32	QSO	0.689	
00 44 59.8	-27 28 20	20.43	-0.58	+0.52	NL	0.238	
00 45 04.5	-30 02 54	18.27	-1.05	+0.25	QSO	2.013	b

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 45 10.0	-29 24 01	18.94	-1.02	+0.45	QSO	1.926	
00 45 18.8	-28 40 14	20.02	-1.15	+0.38	QSO	0.936	d
00 45 22.1	-26 34 51	20.16	-0.82	+0.03	QSO	1.757	
00 45 27.8	-27 42 35	18.18	-0.41	+0.45	QSO	0.232	b,d
00 45 32.9	-26 16 38	19.01	-0.39	-0.02	NL	0.059	
00 45 33.0	-28 55 16	19.71	-0.89	+0.34	QSO	1.183	
00 45 40.6	-25 51 27	19.45	-0.58	+0.35	QSO	2.521	a
00 45 41.0	-28 50 30	18.91	-0.69	+0.22	QSO	2.252	a
00 45 41.4	-28 06 22	18.54	-1.03	+0.34	QSO	1.138	a
00 45 41.6	-27 44 09	19.79	-0.97	+1.16	QSO	0.322	
00 45 44.8	-29 25 57	20.49	-0.71	+0.87	NL	0.204	
00 45 44.8	-27 45 46	19.37	-0.63	+0.38	NL	0.182	
00 45 45.2	-26 06 24	18.08	-0.73	+0.27	QSO	1.242	a
00 45 49.7	-26 04 06	18.89	-0.91	+0.47	QSO	0.486	a
00 45 49.7	-29 40 40	19.26	-0.34	+0.04	QSO	1.087	
00 45 55.8	-29 58 54	19.75	-1.01	+0.65	QSO	0.700	
00 46 01.6	-28 12 40	18.32	-1.05	+0.37	QSO	1.687	a
00 46 02.3	-29 04 54	19.27	-0.51	+0.39	QSO	0.840	c
00 46 09.7	-26 11 36	18.34	-0.50	+1.09	QSO	0.230	e
00 46 13.5	-29 52 22	20.23	-1.22	+0.14	QSO	0.636	
00 46 16.8	-25 45 59	19.78	-0.83	+0.32	QSO	1.018	
00 46 17.8	-28 34 01	17.80	-1.21	+0.43	QSO	0.632	a
00 46 27.4	-29 20 12	20.01	-1.14	+0.37	QSO	1.135	
00 46 34.0	-27 40 43	19.30	-0.86	+0.23	QSO	1.699	
00 46 38.2	-27 48 33	20.08	-0.74	+0.64	NL	0.302	
00 46 38.7	-30 02 08	20.16	-0.63	+0.62	QSO	0.544	
00 46 40.5	-26 09 01	19.48	-0.98	+0.56	QSO	0.435	
00 46 40.6	-25 52 14	20.38	-1.17	+0.80	QSO	1.892	
00 46 45.4	-29 55 55	19.58	-0.71	+0.28	QSO	2.366	
00 46 50.5	-29 14 40	18.12	-1.06	+0.60	QSO	0.781	a
00 46 50.7	-25 41 01	20.34	-1.05	+0.07	QSO	2.166	
00 46 51.5	-26 45 53	19.50	-0.44	+0.47	QSO	2.580	e
00 46 57.8	-27 39 05	20.34	-0.68	+0.38	NL	0.477	
00 46 58.9	-30 11 39	19.17	-1.12	+0.28	QSO	1.081	
00 47 02.4	-27 53 35	19.62	-1.40	+0.75	QSO	1.244	
00 47 06.0	-29 22 54	19.43	-0.61	+0.11	QSO	2.400	
00 47 08.6	-28 49 43	19.87	-1.01	+0.17	QSO	1.653	
00 47 14.9	-29 21 31	19.48	-1.05	+0.32	QSO	2.150	
00 47 15.5	-29 44 44	20.05	-0.66	+0.19	QSO	2.222	
00 47 17.4	-26 14 25	19.53	-0.36	-0.05	QSO	0.781	
00 47 17.6	-26 58 31	19.21	-0.63	+0.48	QSO	0.893	d
00 47 20.7	-28 22 22	20.14	-0.58	+0.96	NL	0.332	
00 47 30.4	-26 33 18	20.40	-0.87	+0.58	QSO	0.970	
00 47 42.0	-26 01 31	19.80	-0.74	+0.71	QSO	1.526	
00 47 43.3	-27 40 05	18.76	-0.52	+0.20	QSO	0.789	
00 47 45.4	-26 33 48	19.95	-0.57	+0.38	QSO	0.414	
00 47 49.5	-27 59 36	18.44	-1.15	+0.40	QSO	2.143	a
00 47 53.4	-26 09 35	19.80	-0.96	+0.43	QSO	0.867	
00 47 54.2	-26 47 53	18.62	-0.79	+0.18	QSO	0.496	a
00 48 02.2	-29 00 46	19.63	-1.48	+0.51	QSO	1.257	
00 48 02.4	-29 47 47	20.25	-1.30	+0.63	QSO	1.251	
00 48 03.2	-28 20 42	20.34	-0.82	+0.30	QSO	1.103	
00 48 07.5	-29 52 34	20.19	-0.83	+0.55	QSO	1.244	
00 48 07.9	-28 18 46	20.04	-0.48	+0.50	QSO	1.322	f
00 48 14.0	-25 57 40	18.69	-0.70	+0.32	QSO	0.780	a
00 48 14.3	-25 55 21	19.31	-1.15	+0.30	QSO	0.825	
00 48 14.4	-27 43 05	19.59	-0.88	+0.41	QSO	0.970	

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 48 16.1	-29 52 37	19.23	-1.59	+0.31	QSO	2.028	a
00 48 16.6	-27 36 38	19.46	-1.00	+0.72	QSO	2.460	
00 48 16.6	-27 40 18	19.76	-0.85	+0.22	QSO	0.675	d
00 48 19.2	-27 56 06	20.04	-0.65	+0.49	QSO	2.146	
00 48 21.0	-29 36 54	20.30	-1.31	+0.62	QSO	2.083	
00 48 21.7	-28 00 09	19.21	-0.15	+0.08	QSO	2.120	d
00 48 21.8	-26 05 39	19.62	-1.35	+0.40	QSO	1.065	
00 48 24.1	-29 01 46	18.28	-0.68	+0.31	QSO	0.783	a
00 48 24.7	-26 01 00	20.08	-1.24	+0.62	QSO	1.317	
00 48 26.3	-29 18 16	18.30	-0.50	+0.14	QSO	0.428	a
00 48 28.1	-29 21 16	19.63	-0.89	+0.27	QSO	1.602	f
00 48 30.4	-29 22 34	20.41	-0.90	+0.50	QSO	1.766	f
00 48 33.6	-28 41 16	19.79	-1.34	+0.68	QSO	1.357	
00 48 34.9	-27 26 04	19.41	-0.66	+0.28	QSO	1.428	
00 48 35.0	-25 45 06	19.26	-1.20	+0.43	QSO	2.082	a
00 48 35.2	-29 49 46	19.72	-0.56	+0.34	QSO	2.439	a
00 48 42.4	-26 08 34	19.12	-0.98	+0.24	QSO	2.249	a
00 48 43.4	-25 46 32	20.10	-1.17	+0.38	QSO	0.904	
00 48 44.7	-29 59 22	19.61	-0.58	+0.49	NL	0.171	
00 48 46.5	-30 09 08	20.47	-1.21	+0.32	QSO	1.530	
00 48 46.9	-28 04 19	17.80	-0.74	+0.50	QSO	0.846	b
00 48 50.5	-29 41 25	19.88	-0.40	+0.26	NL	0.072	
00 48 52.0	-29 07 21	19.77	-0.71	+0.20	QSO	2.370	f
00 48 54.5	-27 24 23	20.45	-0.80	+0.68	QSO	1.268	
00 48 54.9	-29 44 48	19.47	-0.69	+0.19	QSO	2.210	a
00 49 01.1	-28 20 53	18.42	-0.86	+0.37	QSO	2.249	f
00 49 11.3	-26 10 55	20.15	-0.68	+0.33	QSO	2.037	
00 49 13.1	-27 53 11	18.41	-0.72	+0.61	QSO	0.410	b
00 49 14.9	-27 41 48	19.93	-0.98	+0.53	QSO	1.145	
00 49 21.0	-28 07 48	19.66	-1.28	+0.63	QSO	1.148	f
00 49 27.5	-29 31 19	19.15	-0.85	+0.61	QSO	0.601	f
00 49 27.5	-28 12 36	19.49	-0.94	+0.47	QSO	1.145	f
00 49 29.3	-27 14 01	18.99	-0.50	+0.20	QSO	2.484	c
00 49 31.3	-29 18 57	19.83	-1.24	+0.46	QSO	1.631	f
00 49 38.6	-28 00 35	20.20	-0.84	+0.44	QSO	2.479	
00 49 41.4	-28 05 16	20.40	-0.66	+0.21	QSO	1.702	f
00 49 41.4	-29 06 35	19.89	-0.60	+0.45	QSO	0.466	f
00 49 42.2	-28 40 28	18.68	-0.48	+0.12	QSO	0.660	c
00 49 42.8	-26 13 54	19.31	-0.87	+0.08	QSO	0.799	
00 49 43.7	-30 24 15	17.57	-0.79	+0.22	QSO	0.471	b
00 49 44.6	-27 36 12	18.86	-1.19	+0.45	QSO	1.039	f
00 49 46.1	-29 21 41	19.30	-1.10	+0.14	QSO	1.856	f
00 49 46.8	-27 05 12	20.24	-0.64	+1.25	QSO	1.850	
00 49 50.4	-27 49 24	19.77	-0.67	+0.76	QSO	1.007	f
00 49 50.7	-30 26 31	20.43	-0.98	+0.51	QSO	1.339	
00 49 56.5	-27 46 19	19.57	-0.71	+0.50	QSO	0.955	f
00 49 59.1	-29 44 58	18.09	-0.34	+0.14	NL	0.114	b
00 50 01.1	-30 26 17	20.22	-0.80	+0.59	QSO	1.100	
00 50 02.3	-28 15 48	20.24	-1.14	+0.15	QSO	1.892	
00 50 04.7	-26 12 21	20.19	-1.07	+0.78	QSO	1.449	
00 50 04.8	-29 18 20	20.42	-0.76	-0.03	QSO	1.138	
00 50 05.7	-28 04 24	20.11	-1.10	+0.09	QSO	2.152	f
00 50 05.8	-29 05 37	18.39	-0.45	+0.24	QSO	1.605	f
00 50 10.1	-28 07 09	18.91	-0.99	+0.53	QSO	1.730	f
00 50 11.8	-29 03 57	19.89	-0.98	-0.02	QSO	1.619	f
00 50 12.6	-28 45 51	20.12	-0.91	+0.57	QSO	0.727	d
00 50 14.8	-28 24 12	20.09	-0.88	+0.83	QSO	0.551	

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 50 15.7	-26 22 44	18.43	-0.48	+0.70	NL	0.176	b
00 50 18.6	-30 13 58	20.40	-0.26	+0.35	NL	0.075	
00 50 20.4	-28 37 29	19.67	-0.92	+0.80	QSO	0.456	
00 50 23.9	-28 17 40	20.03	-1.20	+0.93	QSO	1.331	f
00 50 26.8	-28 42 12	18.51	-0.94	+0.29	QSO	1.650	b
00 50 28.5	-30 01 01	19.73	-1.43	+0.28	QSO	1.922	a
00 50 33.0	-26 41 28	18.42	-1.32	+0.88	QSO	1.248	a
00 50 35.8	-29 37 03	20.15	-0.99	+0.50	QSO	0.917	f
00 50 36.9	-29 29 13	18.57	-0.54	+0.13	QSO	0.830	f
00 50 37.1	-28 25 27	19.81	-0.41	+0.44	QSO	2.475	f
00 50 38.3	-29 22 36	20.37	-1.40	+0.31	QSO	1.800	f
00 50 40.4	-28 30 23	19.44	-0.61	+0.11	QSO	0.553	f
00 50 45.7	-28 30 36	19.10	-0.98	+0.52	QSO	0.758	f
00 50 47.6	-26 04 56	20.16	-0.48	+0.97	NL	0.258	d
00 50 47.9	-28 50 06	18.38	-0.19	+0.12	NL	0.095	
00 50 49.1	-28 06 29	18.69	-0.87	+0.40	QSO	1.805	a
00 50 50.9	-27 42 06	17.70	-0.65	+0.04	QSO	0.481	a
00 50 58.3	-28 35 47	19.26	-0.81	+0.34	QSO	1.433	
00 51 03.5	-27 45 09	20.04	-0.67	+0.32	QSO	1.178	f
00 51 06.1	-26 08 35	19.39	-1.50	+0.55	QSO	1.262	
00 51 10.1	-26 47 31	19.38	-0.45	+1.16	QSO	1.690	c
00 51 11.7	-25 44 47	19.45	-1.32	+1.23	QSO	1.390	
00 51 12.2	-27 15 32	19.32	-1.37	+0.61	QSO	1.200	
00 51 13.5	-29 19 40	19.28	-1.20	+0.64	QSO	1.486	f
00 51 13.9	-27 33 41	18.87	-0.84	+0.49	QSO	1.611	f
00 51 17.4	-29 16 43	20.34	-0.73	+0.21	QSO	1.107	f
00 51 21.6	-28 39 58	18.33	-0.88	+0.47	QSO	1.574	f
00 51 23.3	-30 24 30	19.90	-0.58	+0.95	QSO	2.116	
00 51 29.1	-29 17 13	19.11	-0.50	+0.66	QSO	1.479	f
00 51 34.4	-28 36 26	19.08	-0.17	+0.14	QSO	0.601	
00 51 34.7	-27 55 54	20.44	-0.68	+0.81	NL	0.409	
00 51 36.3	-30 13 57	19.96	-0.70	+0.41	QSO	1.767	
00 51 36.4	-29 24 16	20.47	-0.49	+0.76	QSO	1.550	f
00 51 38.1	-27 26 26	18.87	-1.00	+0.19	QSO	0.689	f
00 51 39.5	-28 46 48	19.43	-1.60	+0.79	QSO	1.338	f
00 51 42.5	-28 01 13	18.51	-0.47	+0.65	QSO	1.504	b
00 51 46.7	-28 11 52	18.83	-0.90	+0.29	QSO	2.244	a
00 51 48.6	-30 12 03	18.69	-1.13	+0.67	QSO	1.141	a
00 51 51.4	-29 18 36	19.97	-0.88	+0.07	QSO	1.987	f
00 51 52.3	-28 22 28	19.23	-0.64	+0.25	QSO	2.133	f
00 51 53.4	-26 05 14	18.04	-0.55	+0.08	QSO	0.624	a
00 51 58.3	-29 47 04	20.36	-0.70	+0.52	NL	0.216	
00 52 00.2	-30 21 21	19.87	-1.03	+0.60	QSO	1.464	
00 52 00.7	-30 20 54	17.55	-0.80	+0.24	QSO	0.993	a
00 52 03.8	-28 09 24	19.74	-0.46	+0.52	QSO	0.838	
00 52 09.1	-27 40 57	20.25	-0.53	+0.63	NL	0.156	
00 52 13.7	-29 24 15	19.18	-0.94	+0.51	QSO	1.424	f
00 52 16.0	-28 32 32	20.23	-0.79	+0.39	QSO	2.181	f
00 52 16.7	-26 15 57	19.20	-0.95	+0.30	QSO	0.850	
00 52 18.0	-28 47 35	19.24	-0.92	+0.33	QSO	0.639	f
00 52 18.5	-26 01 56	20.05	-0.98	+0.10	QSO	1.925	
00 52 20.8	-28 00 17	20.34	-1.43	+0.34	QSO	1.725	
00 52 21.9	-28 48 11	20.18	-0.75	+0.21	QSO	2.097	f
00 52 22.6	-29 03 30	19.93	-0.47	+0.96	QSO	1.053	
00 52 26.9	-29 45 50	18.58	-0.35	+0.09	QSO	0.760	c
00 52 27.9	-30 08 25	19.80	-0.98	+0.20	QSO	0.950	g
00 52 33.7	-28 30 46	19.42	-0.86	+0.28	QSO	0.779	f

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 52 35.8	-27 12 39	20.05	-0.75	+0.14	QSO	1.264	
00 52 38.4	-28 47 58	19.80	-0.64	+0.58	QSO	0.777	
00 52 38.8	-28 23 48	19.36	-0.69	+0.37	QSO	2.350	f
00 52 40.9	-28 56 49	18.33	-0.54	+0.34	QSO	0.602	a
00 52 41.1	-30 30 37	19.17	-1.39	+0.41	QSO	1.146	
00 52 41.1	-25 50 36	18.51	-0.72	-0.07	QSO	0.854	a
00 52 42.6	-30 11 39	19.96	-0.71	+1.08	QSO	1.380	g
00 52 44.5	-29 02 32	19.08	-0.20	+0.22	QSO	0.840	c
00 52 46.5	-30 06 20	19.81	-1.02	+0.09	QSO	1.788	
00 52 51.3	-28 53 35	18.65	-0.51	+0.24	QSO	0.634	a
00 52 55.1	-30 15 50	19.95	-0.90	+0.41	QSO	2.117	
00 52 57.6	-29 42 05	20.17	-0.37	+0.39	QSO	2.367	
00 53 00.3	-28 58 15	20.02	-1.14	+0.62	QSO	0.940	d
00 53 02.0	-25 53 05	19.94	-0.62	+0.18	QSO	2.230	c
00 53 06.8	-27 24 10	19.69	-1.03	+0.51	QSO	0.433	d
00 53 08.0	-27 38 59	19.64	-0.71	-0.06	QSO	2.107	
00 53 11.3	-26 39 01	18.44	-0.80	+0.46	QSO	0.808	a
00 53 15.2	-29 24 42	19.70	-0.89	+0.56	QSO	1.331	f
00 53 19.1	-26 56 45	19.67	-1.09	+0.43	QSO	1.882	
00 53 19.5	-29 21 53	19.58	-1.30	+0.31	QSO	2.029	f
00 53 19.9	-28 13 49	20.44	-1.68	+0.34	QSO	1.860	a
00 53 20.3	-30 16 14	19.31	-0.31	+0.27	QSO	2.440	c
00 53 24.1	-28 45 24	19.25	-0.65	+0.33	QSO	1.703	a
00 53 30.9	-28 36 27	18.78	-0.94	+0.48	QSO	1.920	f
00 53 31.7	-29 00 12	20.13	-1.02	+0.76	QSO	1.926	
00 53 33.8	-28 36 50	20.37	-1.42	+0.50	QSO	1.306	f
00 53 34.7	-29 37 57	19.85	-0.58	+0.83	NL	0.306	
00 53 37.0	-28 43 13	19.35	-1.43	+0.33	QSO	1.933	f
00 53 37.0	-28 30 13	19.52	-0.62	+0.42	QSO	1.029	f
00 53 38.5	-27 09 10	18.02	-0.43	+0.28	QSO	1.039	a
00 53 39.4	-30 28 24	19.11	-1.25	+0.33	QSO	1.190	
00 53 41.6	-29 04 58	20.34	-0.51	+0.36	QSO	2.222	
00 53 43.5	-29 24 15	18.97	-0.54	+0.75	NL	0.271	f
00 53 45.5	-27 19 12	20.03	-0.63	+0.33	QSO	0.536	
00 53 47.1	-28 13 25	18.44	-0.36	+0.06	QSO	0.725	a
00 53 48.4	-30 07 14	19.75	-0.94	+0.31	QSO	1.040	g
00 53 52.6	-29 30 16	19.91	-1.06	+0.31	QSO	1.969	f
00 53 59.1	-29 52 07	19.87	-1.02	+0.22	QSO	1.605	
00 54 03.6	-29 17 19	18.76	-0.93	+0.38	QSO	1.801	f
00 54 06.3	-27 33 36	18.96	-1.12	+0.67	QSO	1.260	a
00 54 09.6	-28 32 36	20.19	-0.55	+0.21	QSO	1.928	
00 54 15.9	-26 08 57	20.13	-0.68	+0.55	QSO	0.604	
00 54 16.6	-29 34 35	19.67	-0.97	+0.78	QSO	1.417	
00 54 18.8	-26 49 57	20.26	-1.07	+0.55	QSO	1.606	
00 54 19.6	-26 19 05	20.40	-0.81	+0.15	QSO	1.084	
00 54 23.6	-29 53 53	19.35	-1.36	+0.51	QSO	1.660	
00 54 25.8	-28 26 08	19.58	-0.80	+0.50	QSO	0.772	
00 54 26.2	-25 53 32	19.40	-1.21	+0.52	QSO	0.785	
00 54 27.4	-29 38 34	19.97	-0.94	+0.56	QSO	1.418	
00 54 29.2	-28 07 29	20.48	-1.29	+0.72	QSO	1.849	
00 54 30.3	-29 04 17	20.46	-0.57	+0.34	NL	0.229	
00 54 30.3	-29 36 29	18.92	-0.78	+0.46	QSO	1.610	
00 54 32.2	-30 23 56	19.83	-0.41	+0.41	NL	0.157	
00 54 33.0	-29 34 06	19.42	-0.89	+0.24	QSO	2.050	
00 54 34.6	-27 18 42	20.30	-0.62	+0.62	NL	0.291	
00 54 35.7	-27 44 13	19.32	-0.82	+0.41	QSO	0.348	f
00 54 38.0	-30 05 20	19.79	-0.99	+0.28	QSO	1.582	

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 54 51.9	-26 43 16	19.62	-0.52	+0.92	NL	0.241	
00 54 56.2	-30 07 18	20.36	-0.99	+1.21	QSO	0.366	
00 54 57.9	-29 40 02	20.39	-0.97	+0.48	QSO	1.428	
00 54 59.2	-27 48 13	19.11	-0.85	+0.53	QSO	1.209	f
00 55 01.0	-26 19 48	19.71	-0.56	+0.65	NL	0.131	
00 55 06.4	-27 21 50	19.00	-0.64	+0.46	QSO	0.898	
00 55 09.1	-27 06 14	20.09	-0.73	+0.48	QSO	1.098	
00 55 09.6	-27 44 40	18.41	-0.40	+0.31	QSO	2.174	f
00 55 11.9	-29 57 33	19.67	-0.89	+0.36	QSO	1.463	
00 55 14.3	-25 59 05	18.04	-0.67	+0.23	QSO	0.584	a
00 55 15.2	-28 39 23	19.51	-0.84	+0.11	QSO	1.366	f
00 55 17.2	-27 20 26	19.41	-0.95	+0.61	QSO	1.318	
00 55 19.3	-27 19 44	20.47	-0.57	+0.58	NL	0.198	
00 55 20.6	-27 31 09	20.48	-0.60	+0.08	QSO	1.640	
00 55 26.2	-27 34 17	19.11	-0.20	+0.21	NL	0.186	
00 55 27.4	-26 47 53	19.51	-0.50	+1.02	NL	0.181	
00 55 28.9	-26 23 27	20.46	-1.54	+0.27	QSO	1.959	
00 55 28.9	-27 43 12	20.48	-1.13	+0.82	QSO	0.907	f
00 55 31.3	-26 12 25	19.94	-0.37	+0.39	QSO	0.793	
00 55 32.5	-26 59 26	18.92	-0.60	+1.25	QSO	3.662	a
00 55 34.7	-28 03 45	19.18	-0.76	+0.48	QSO	1.650	f
00 55 36.3	-27 50 41	20.17	-0.59	+0.93	NL	0.274	
00 55 38.4	-28 28 23	18.71	-0.51	+0.19	QSO	0.648	a
00 55 40.7	-28 06 16	20.20	-0.65	+1.11	QSO	0.486	
00 55 42.9	-28 50 14	19.22	-0.74	+0.38	QSO	1.276	f
00 55 43.4	-29 49 00	18.58	-0.79	+0.34	QSO	0.668	a
00 55 45.6	-26 54 12	20.08	-0.31	+0.28	QSO	2.390	
00 55 45.9	-28 24 29	20.46	-0.90	+0.12	QSO	0.614	
00 55 45.9	-29 59 09	19.97	-1.10	+0.19	QSO	0.613	
00 55 46.5	-27 36 32	19.95	-1.21	+0.33	QSO	1.759	
00 55 53.0	-27 44 60	19.77	-0.87	+0.31	QSO	1.606	f
00 55 54.1	-29 58 39	20.30	-0.58	+0.19	QSO	2.202	
00 55 54.9	-28 21 55	20.20	-1.25	+0.67	QSO	1.418	
00 55 57.3	-26 38 55	19.88	-0.91	+0.19	QSO	0.740	
00 55 59.4	-27 42 17	20.41	-0.90	+0.63	NL	0.211	f
00 56 03.3	-27 16 15	20.05	-1.31	+0.51	QSO	1.614	
00 56 03.4	-25 58 12	19.30	-0.86	+0.51	QSO	0.930	
00 56 05.5	-29 57 34	20.33	-0.79	+0.58	QSO	0.652	
00 56 06.5	-27 54 08	20.12	-0.36	+0.43	QSO	2.861	f
00 56 07.9	-30 23 18	19.91	-0.30	+0.03	QSO	2.100	c
00 56 11.3	-26 09 11	19.01	-0.69	+0.57	QSO	1.667	
00 56 11.5	-26 39 49	20.34	-0.95	+0.65	QSO	1.384	
00 56 11.8	-29 18 34	18.97	-0.38	+0.08	QSO	0.910	c
00 56 12.5	-28 43 28	19.94	-0.75	+0.45	QSO	0.828	f
00 56 13.3	-27 03 27	19.31	-0.80	+0.36	QSO	0.918	d
00 56 16.2	-30 03 17	20.17	-0.43	+0.41	QSO	2.202	
00 56 16.4	-30 18 14	19.39	-0.67	+0.38	QSO	1.362	
00 56 18.0	-28 41 45	20.41	-0.58	+0.42	QSO	0.965	f
00 56 18.2	-28 45 55	19.26	-0.56	+0.27	QSO	1.740	f
00 56 20.9	-29 15 35	18.56	-1.01	+0.53	QSO	1.255	a
00 56 21.6	-28 47 47	20.49	-1.07	+0.55	QSO	0.800	
00 56 21.7	-29 11 24	19.31	-0.91	+0.86	QSO	1.408	
00 56 22.2	-29 48 05	20.42	-0.51	+0.55	NL	0.199	
00 56 27.4	-27 45 43	20.42	-0.92	+0.62	QSO	0.560	f
00 56 27.6	-28 32 43	20.02	-0.53	+0.98	NL	0.163	
00 56 31.4	-26 29 29	19.25	-0.11	+0.22	QSO	0.829	
00 56 32.7	-27 56 35	18.61	-0.15	+0.17	NL	0.095	

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 56 34.6	-29 05 32	18.97	-0.79	+0.49	QSO	1.341	f
00 56 35.2	-26 19 49	19.02	-1.07	+0.65	QSO	1.116	
00 56 35.4	-27 00 46	20.32	-0.62	+0.21	QSO	2.284	
00 56 41.4	-28 43 15	17.69	-0.84	+0.37	QSO	0.934	a
00 56 43.9	-29 49 36	19.66	-0.80	+0.57	QSO	1.282	
00 56 46.8	-27 38 45	19.01	-0.44	+0.13	QSO	1.618	
00 56 53.3	-28 17 38	20.04	-0.62	+1.16	NL	0.277	
00 56 53.4	-26 56 35	20.21	-0.51	+1.02	NL	0.333	
00 56 58.1	-29 48 20	18.09	-0.11	+0.14	QSO	0.351	a
00 56 58.4	-25 59 06	19.75	-1.52	+0.46	QSO	1.085	
00 57 01.9	-28 35 43	19.34	-0.77	+0.38	QSO	0.662	f
00 57 03.6	-25 45 34	20.42	-0.91	+0.36	QSO	1.228	
00 57 04.6	-27 59 48	19.90	-1.46	+0.69	QSO	1.195	f
00 57 07.2	-27 41 40	20.34	-1.29	+0.44	QSO	1.684	
00 57 10.2	-30 20 06	18.38	-0.54	+0.29	QSO	0.394	b
00 57 12.2	-29 21 18	18.98	-1.00	+0.52	QSO	0.955	
00 57 14.4	-26 39 34	19.97	-1.62	+0.67	QSO	1.042	
00 57 16.6	-25 41 20	18.71	-1.09	+0.53	QSO	1.262	
00 57 19.1	-27 54 40	19.24	-0.93	+0.32	QSO	1.698	
00 57 23.4	-25 55 41	18.94	-1.05	+0.45	QSO	0.315	a
00 57 26.1	-29 29 50	20.37	-1.16	+0.21	QSO	0.681	
00 57 27.9	-29 55 02	20.29	-1.07	+0.45	QSO	0.656	
00 57 29.1	-26 23 29	19.43	-0.53	+0.87	NL	0.132	
00 57 30.2	-26 30 22	18.58	-0.79	+0.40	QSO	1.042	a
00 57 30.6	-28 48 51	19.77	-0.57	+0.53	QSO	2.867	f
00 57 36.6	-29 03 28	20.22	-0.42	+0.48	QSO	2.380	
00 57 36.8	-29 20 50	19.40	-0.85	+0.37	QSO	1.072	
00 57 40.1	-29 38 40	19.36	-0.36	+0.27	QSO	0.512	
00 57 40.7	-29 08 16	18.32	-0.87	+0.37	QSO	0.489	a
00 57 42.6	-26 48 28	19.91	-1.22	+0.71	QSO	1.304	
00 57 44.4	-27 08 06	20.19	-1.03	+0.90	QSO	1.189	
00 57 44.8	-27 58 28	19.35	-0.98	+0.53	QSO	1.134	
00 57 47.4	-28 22 41	19.03	-0.12	+0.18	NL	0.053	
00 57 48.1	-30 30 44	20.30	-0.91	+0.25	QSO	1.152	
00 57 51.0	-30 27 25	20.46	-1.13	+0.72	QSO	1.228	
00 57 56.7	-27 29 46	18.43	-0.92	+0.58	QSO	1.203	a
00 57 57.8	-28 37 08	19.26	-0.77	+0.38	QSO	0.669	d
00 57 59.5	-30 16 06	20.50	-1.25	+0.32	QSO	1.446	
00 57 59.5	-30 15 34	20.50	-0.94	+0.42	QSO	0.774	d
00 58 05.2	-26 18 28	18.06	-0.67	+0.27	QSO	0.792	b
00 58 06.2	-26 04 44	18.50	-0.69	+0.48	QSO	2.472	a
00 58 14.0	-25 54 36	16.73	-0.17	+0.17	NL	0.158	a
00 58 14.5	-28 08 03	20.47	-0.24	+0.02	QSO	0.308	
00 58 20.2	-26 05 18	19.72	-0.93	+0.75	QSO	1.790	
00 58 21.0	-25 52 48	19.86	-0.87	+0.53	QSO	1.712	
00 58 22.4	-28 21 24	19.53	-0.73	+0.91	NL	0.310	
00 58 25.9	-29 31 45	19.16	-1.04	+0.37	QSO	0.872	
00 58 26.0	-30 17 35	20.29	-0.72	+0.63	NL	0.241	
00 58 27.2	-30 08 24	18.78	-1.20	+0.46	QSO	1.263	
00 58 27.8	-27 58 44	19.80	-0.56	+0.91	NL	0.199	
00 58 28.9	-29 51 44	19.10	-0.70	+0.15	QSO	0.915	
00 58 28.9	-29 56 22	19.68	-0.90	+0.41	QSO	1.090	
00 58 34.6	-29 19 02	18.27	-0.73	+0.40	QSO	1.184	a
00 58 35.7	-29 07 24	18.00	-0.77	+0.32	QSO	0.866	a
00 58 36.2	-29 39 27	19.33	-1.09	+0.39	QSO	0.964	
00 58 37.9	-26 26 60	20.03	-0.89	+0.63	QSO	2.535	
00 58 39.8	-26 55 13	19.81	-0.80	+0.59	QSO	2.350	a

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
00 58 41.4	-28 31 01	18.62	-0.69	+0.37	QSO	1.001	a
00 58 42.6	-29 24 08	19.95	-0.09	+0.11	QSO	1.620	c
00 58 44.4	-25 57 53	19.22	-0.65	+0.58	QSO	1.704	
00 58 44.5	-25 41 27	19.90	-1.61	+0.73	QSO	0.667	d
00 58 54.7	-29 40 05	19.36	-0.64	+0.66	QSO	0.721	
00 58 55.7	-29 58 46	19.63	-0.72	+0.51	QSO	1.305	
00 58 57.2	-28 56 58	20.39	-0.93	+1.42	QSO	0.322	d
00 58 57.6	-27 05 42	19.52	-1.13	+0.52	QSO	1.889	a
00 58 58.2	-28 23 37	20.36	-0.49	+0.34	QSO	1.554	
00 59 00.6	-28 13 44	19.80	-1.07	+0.54	QSO	1.160	
00 59 01.7	-30 07 56	20.13	-0.52	+0.42	QSO	2.780	
00 59 05.3	-27 22 28	20.08	-0.61	+0.20	QSO	1.617	
00 59 06.8	-30 25 53	18.89	-0.12	+0.21	QSO	2.170	c
00 59 08.2	-27 29 03	17.78	-0.73	+0.77	QSO	0.188	a
00 59 10.7	-28 53 39	18.14	-0.29	+0.11	QSO	0.620	c
00 59 12.5	-28 45 37	19.92	-1.18	+0.39	QSO	0.760	
00 59 14.3	-26 55 10	19.28	-1.47	+0.63	QSO	1.367	
00 59 17.6	-28 29 55	20.05	-1.02	+0.07	QSO	1.676	
00 59 25.9	-30 19 03	19.96	-0.94	+0.71	QSO	1.350	d
00 59 26.1	-29 09 22	20.38	-0.81	+0.74	QSO	2.647	
00 59 27.8	-26 25 06	18.51	-0.91	+0.13	QSO	2.109	a
00 59 28.4	-29 11 22	20.25	-0.28	+0.32	NL	0.197	
00 59 31.4	-30 08 55	19.50	-0.81	+0.37	QSO	1.505	
00 59 35.0	-30 26 33	19.31	-0.87	+0.32	QSO	1.262	
00 59 37.3	-30 34 36	16.92	-0.51	+0.28	QSO	1.033	a
00 59 38.7	-27 02 14	20.45	-1.19	+0.04	QSO	1.970	e
00 59 39.4	-27 02 07	18.91	-1.13	+0.34	QSO	0.940	e
00 59 42.5	-26 10 50	19.51	-1.09	+0.30	QSO	1.397	
00 59 47.2	-26 15 58	19.77	-0.74	+0.36	QSO	0.778	
00 59 49.8	-29 29 00	19.35	-0.15	+0.32	NL	0.058	
00 59 50.4	-29 46 20	17.73	-0.85	+0.28	QSO	1.076	a
00 59 54.0	-29 40 34	20.19	-1.04	+0.10	QSO	2.143	
00 59 57.2	-26 57 53	18.62	-0.71	+0.27	QSO	2.266	a
00 59 59.3	-27 42 17	20.35	-0.85	+0.51	QSO	0.463	
00 59 60.0	-26 06 54	20.20	-1.18	+0.65	QSO	1.694	
01 00 02.1	-28 28 14	20.36	-1.04	+0.42	QSO	1.730	
01 00 02.7	-30 32 01	19.78	-0.84	+0.25	QSO	0.929	
01 00 03.1	-26 37 25	19.78	-0.79	+0.34	NL	0.171	
01 00 10.2	-28 33 35	19.64	-1.17	+0.68	QSO	1.350	
01 00 11.2	-28 11 32	20.19	-0.27	+0.26	QSO	1.112	
01 00 15.9	-28 43 04	20.32	-0.76	+0.82	NL	0.399	
01 00 16.0	-29 10 21	18.95	-0.58	+0.29	QSO	2.380	a
01 00 24.2	-27 13 33	19.76	-0.96	+0.30	QSO	1.313	
01 00 25.4	-28 21 33	19.20	-0.17	+0.30	QSO	2.640	c
01 00 26.0	-30 01 58	20.04	-0.80	+0.51	QSO	0.500	
01 00 27.6	-28 09 10	18.04	-0.49	+0.86	QSO	1.768	a
01 00 29.7	-28 47 60	19.00	-0.24	+0.17	NL	0.097	
01 00 31.7	-27 02 42	17.78	-1.06	+0.32	QSO	1.597	a
01 00 33.1	-28 09 45	19.38	-1.04	+0.42	QSO	1.800	a
01 00 33.2	-26 10 31	18.81	-0.94	+0.46	QSO	2.540	a
01 00 33.3	-29 12 50	20.22	-0.41	+0.39	QSO	2.760	c
01 00 35.3	-27 39 49	20.23	-1.82	+1.12	QSO	1.418	a
01 00 37.6	-26 29 39	20.27	-1.11	+0.41	QSO	1.038	
01 00 40.5	-29 55 52	20.21	-0.14	+0.26	QSO	2.681	
01 00 42.5	-30 37 07	18.74	-0.40	+0.26	QSO	1.002	
01 00 46.3	-26 29 46	20.24	-1.01	+0.46	QSO	1.305	
01 00 47.1	-30 30 54	18.66	-0.84	+0.40	QSO	0.940	b

Table 3. continued

α (1950)	δ (1950)	B_j	$U - B_j$	$B_j - R$	ID	z	Comm
01 00 50.8	-30 23 58	19.62	-1.27	+0.41	QSO	0.946	
01 00 50.9	-28 57 29	19.09	-1.01	+0.44	QSO	1.226	a
01 00 52.1	-26 49 56	17.34	-0.19	+0.13	NL	0.156	
01 00 59.5	-26 22 31	20.11	-1.45	+0.52	QSO	1.022	
01 01 01.9	-27 08 50	18.50	-0.76	-0.00	QSO	0.558	a
01 01 03.3	-28 01 30	20.13	-0.64	+0.38	QSO	1.564	
01 01 03.6	-26 35 33	20.32	-1.00	+0.43	QSO	2.091	
01 01 04.3	-29 58 55	18.81	-0.25	-0.19	QSO	0.760	c
01 01 05.0	-26 46 51	19.74	-1.02	+0.25	QSO	1.795	
01 01 06.1	-25 43 57	20.23	-0.36	+0.30	QSO	1.765	
01 01 06.2	-29 57 22	20.20	-1.05	+0.15	QSO	1.802	
01 01 08.3	-25 48 31	18.58	-1.00	+0.17	QSO	1.973	a
01 01 11.5	-26 50 31	20.05	-1.09	+0.29	QSO	1.815	
01 01 13.6	-29 58 39	20.07	-0.58	+0.37	NL	0.107	
01 01 14.5	-29 32 05	19.31	-0.75	+0.74	QSO	0.326	
01 01 27.3	-30 22 28	19.22	-0.90	+0.46	QSO	1.326	
01 01 27.8	-28 55 55	19.93	-0.73	+0.65	QSO	1.036	
01 01 30.7	-30 29 36	20.25	-1.67	-0.11	QSO	0.777	
01 01 30.7	-26 00 12	19.29	-0.68	+0.50	QSO	0.449	
01 01 41.7	-25 48 09	19.67	-0.38	+0.30	QSO	1.536	
01 01 53.6	-30 20 55	20.33	-0.70	+0.69	QSO	0.763	
01 02 07.4	-30 16 46	19.04	-0.54	+0.83	NL	0.177	
01 02 16.7	-27 13 11	17.33	-0.68	+0.38	QSO	0.780	a
01 02 38.8	-28 42 30	18.65	-0.92	+0.53	QSO	1.375	b
01 02 39.5	-29 22 22	19.08	-0.57	+0.47	QSO	2.440	c
01 02 43.2	-30 12 01	17.78	-0.60	+0.15	QSO	0.838	a
01 02 51.2	-30 04 31	20.37	-0.32	+0.10	QSO	2.359	
01 02 53.7	-29 03 32	19.00	-0.72	+0.47	QSO	1.540	a
01 02 59.5	-30 14 53	18.24	-0.62	+0.09	QSO	0.533	b

^a Veron-Cetty & Veron (1998)^b Cristiani et al. (1995)^c Warren et al. (1991a)^d Uncertain redshift^e Hook et al. (1994)^f Boyle et al. (1990)^g Spectrum missing in Fig. 1




Fig. 1. Spectra of all the new emission line objects identified in the survey.





Fig. 1. continued






Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued




Fig. 1. continued

Acknowledgements. It is a pleasure to thank the enthusiastic support of the COSMOS and UKST staff.

References

- Andreani P., Cristiani S., 1992, ApJ 298, L13
 Beard S.M., MacGillivray H.T., Thanisch P.F., 1990, MNRAS 247, 311
 Bellenger R., Dreux M., Felenbok P., et al., 1991, The Messenger 65, 54
 Blair M., Gilmore G., 1982, PASP 94, 742
 Boyle B.J., Fong R., Shanks T., Peterson B.A., 1990, MNRAS 243, 1
 Campusano L.E., 1991, A&A 259, 9
 Cristiani S., La Franca F., Andreani P., et al., 1995, A&AS 112, 347
 Croom S.M., Shanks T., Boyle B.J., et al., 1998, in Evolution of Large Scale Structure (Garching 1998), astro-ph/9810127
 da Costa L.N., 1998, in "Wide Field Surveys in Cosmology", eds. S. Colombi, Y. Mellier, astro-ph/9809209
 Goldschmidt P., Miller L., 1998, MNRAS 293, 107
 Goldschmidt P., Miller L., La Franca F., Cristiani S., 1992, MNRAS 256, 65p
 Hawkins M.R.S., 1991, IAU Commission 9 - Newsletter 1, 23
 Hawkins M.R.S., Bessell M.S., 1988, MNRAS 234, 177
 Hewett P.C., Foltz C.G., Chaffe F.H., 1995, AJ 109, 1498
 Hook I.M., McMahon R.G., Boyle B.J., Irwin M.J., 1994, MNRAS 268, 305
 La Franca F., Cristiani S., 1997, AJ 113, 1517
 La Franca F., Cristiani S., Barbieri C., 1992, AJ 103, 1062
 La Franca F., Andreani P., Cristiani S., 1998, ApJ 497, 529
 Lissandrini C., Cristiani S., La Franca F., 1994, PASP 106, 1157
 Nonino M., Bertin E., Da Costa L., et al., 1999, A&AS 137, 43
 MacGillivray H.T., Stobie R.S., 1984, Vistas Astron. 27, 433
 Prandoni I., Wichmann R., da Costa L., et al., 1999, A&A 345, 448
 Schmidt M., Green R.G., 1983, ApJ 269, 352
 Veron-Cetty M.P., Veron P., 1998, ESO Sci. Rep. 18, 1
 Warren S.J., Hewett P.C., Irwin M.J., Osmer P.S., 1991a, ApJS 76, 1
 Warren S.J., Hewett P.C., Osmer P.S., 1991b, ApJS 76, 23
 Zaggia S., Hook I., Mendez R., et al., 1999, A&AS 137, 75